

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/279338082>

Carbon sequestration by a few marine algae: Observation and projection

Article · January 2009

CITATIONS

69

READS

511

3 authors, including:



Kaladharan P

Central Marine Fisheries Research Institute

193 PUBLICATIONS 775 CITATIONS

[SEE PROFILE](#)



Veena Shettigar

Central Marine Fisheries Research Institute

27 PUBLICATIONS 238 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Marine macrophytes [View project](#)



Marine Macrophytes in India- Resource dynamics & Ecosystem services [View project](#)



Short Communication

†Carbon sequestration by a few marine algae: observation and projection

¹*P. Kaladharan, ¹S.Veena and ²E. Vivekanandan

¹Visakhapatnam Regional Centre of Central Marine Fisheries Research Institute, Visakhapatnam 530 003, India. *E-mail: kaladharanep@gmail.com

²Central Marine Fisheries Research Institute, Cochin-682018, India.

Abstract

CO₂ sequestration by the marine planktonic microalgae *Nannochloropsis salina* and *Isochrysis galbana* as well as macroforms *Gracilaria corticata*, *Sargassum polycystum* and *Ulva lactuca* was estimated under laboratory conditions. The green seaweed *U. lactuca* registered 100% utilization of CO₂ towards carbon fixation from the ambient water up to 15 mg/l and beyond that it declined to 60%. The microalgae were able to utilize 27.7% of dissolved CO₂ at 15 mg/l, but did not show any effect either for carbon fixation or for emission at lower and higher levels. Gross primary productivity of these algae were also not affected by increase in the CO₂ levels. It is estimated that the seaweed biomass along the Indian coast is capable of utilizing 9052 tCO₂/d against emission of 365 tCO₂/d indicating a net carbon credit of 8687 t/d.

Keywords: Carbon sequestration, microalgae, seaweeds, carbon credits

Introduction

Increase of carbon dioxide besides methane and oxides of nitrogen in the atmosphere is leading to climate change. These greenhouse gases (GHGs) cause depletion of ozone layer protecting the atmosphere against UV radiation, thereby warming the atmosphere. The average concentration of CO₂ increased from 315 ppm in 1960 to 380 ppm in 2007 (IPCC, 2007). There has been a 35% increase in CO₂ emission worldwide since 1990.

Carbon fixation by photoautotrophic algae has the potential to diminish the release of CO₂ into the atmosphere and in helping to alleviate the trend toward global warming. Primary producers of coastal and marine ecosystems such as phytoplankton, seaweed and seagrass are excellent carbon sequestering agents than their terrestrial counterparts (Zou, 2005). In this communication, an attempt is made to assess the carbon fixing efficiency and respiratory emission rate of two marine planktonic algae and three seaweeds in varying levels of

dissolved CO₂ in relation to the proposed rise in the level of dissolved CO₂ in coastal waters. We also attempt to project a quantitative estimate of carbon sequestering capacity of the seaweeds that are distributed along the Indian coast based on our observation of CO₂ assimilation ability of various marine algal species.

Material and methods

Marine algal samples: Samples of *Gracilaria corticata* J. Agardh, *Sargassum polycystum* C. Agardh and *Ulva lactuca* Linn. were collected from the intertidal rocky beach adjoining Thotlakonda (17°49'18.75"N lat. and 83°24'58.06"E long.) near Visakhapatnam and brought to the laboratory. The samples were washed and the thalli were cleaned with excess water to free epiphytes, organisms and adhering sand particles. They were acclimatized overnight in large tanks containing filtered (0.45μ) seawater of 32 ppt. Samples of microalgae *Nannochloropsis salina* Hibberd and *Isochrysis galbana* Parke were taken from the stocks at their

†Presented in the International Symposium "Marine Ecosystem- Challenges and Opportunities (MECOS 09)" organized by the Marine Biological Association of India during February 9-12, 2009 at Kochi.

exponential growth phase, being maintained at the algal feed culture laboratory of Visakhapatnam Regional Centre of CMFRI for the last ten years. The concentration of cells during the experiment was 1.25×10^6 /ml for *Nannochloropsis salina* and 2.0×10^6 /ml for *Isochrysis galbana*.

Enrichment of CO₂: Seawater collected from the seaweed collection site was brought to the laboratory and filtered through 0.45µ GFC (Millipore) to remove plankton and algal spores. Prior to the experiment, the initial CO₂ concentration was determined. Later, CO₂ was dissolved into the seawater taken in separate glass containers at the rate of 5, 10, 15 and 25 mg/l from a CO₂ cylinder dispensed through a tabletop sodamaker. The initial level of dissolved CO₂ in each set of airtight container was ascertained titrimetrically according to Dye (1958) and maintained at the required level.

Determination of CO₂ utilization, emission and primary production: Cleaned thallii of seaweeds were weighed (0.5 g) accurately and incubated with 300 ml of CO₂ enriched seawater (5-25 mg/l) after determining the initial O₂ and CO₂ in airtight, thick and clear polythene bags in light and dark condition under a column of water (40-50 cm) for two hours. After the incubation in light/ dark, 100 ml water samples from each bag were drawn gently using a large hypodermic syringe into separate glass bottles (100 ml) and subjected to CO₂ determination (Dye, 1958) and another 100 ml for determining dissolved oxygen (Gaarder and Gran, 1927) using Winkler's reagent.

Similarly 10 ml each of stock cultures of *Nannochloropsis salina* and *Isochrysis galbana* were treated with 290 ml of CO₂ enriched (0-25 mg/l) seawater. CO₂ utilized for carbon fixation (light incubation), CO₂ emission (dark incubation) and

their gross primary productivity (GPP) in various levels of dissolved CO₂ were determined. GPP was calculated for each species in different CO₂ concentrations from their oxygen production values multiplied with the factor 0.536/PQ, where PQ was 1.25 (Westlake, 1963). Triplicate polythene bags for each species and each CO₂ treatment and control (without seaweeds as well as 0 mg/l of CO₂) were maintained to get average values.

Results

Carbon utilization and emission: Four levels of dissolved CO₂ in seawater were tested to assess the rate of carbon utilization in light and its emission in dark during respiration by the macro and microalgae and the results are presented in Table 1. *Gracilaria corticata* (red alga) and *Sargassum polycystum* (brown alga) utilized 100% of the dissolved CO₂ for photosynthesis when the ambient level was 5 mg/l and beyond that, the CO₂ utilization rate showed steep decline. The emission rate was nil when the ambient CO₂ levels were increased from 5 to 25 mg/l, except at 25 mg/l concentration, where 20% increase in emission over the control was observed in the case of *G. corticata*. However, the green alga *Ulva lactuca* registered 100% utilization of CO₂ for carbon fixation from the ambient water upto a level of 15 mg/l and beyond that level it declined to 60%. The microalgae were able to utilize 27.7% of dissolved CO₂ when the ambient level was 15 mg/l, but at lower and higher levels it did not show any effect either for carbon fixation or for emission.

As shown in Table 2 the effect of increasing levels of dissolved CO₂ on the GPP of *Sargassum polycystum* at elevated levels of CO₂ in ambient water gradually increased from 124% to 209% over

Table 1. CO₂ fixed (% over initial level) in light (photosynthesis) or emitted (% over initial level) in dark (respiration) by some marine algae observed under different levels of dissolved CO₂ in closed system

Species	CO ₂ level (mg/l)									
	0		5		10		15		25	
	Light	Dark	Light	Dark	Light	Dark	Light	Dark	Light	Dark
<i>Gracilaria corticata</i>	0	0	100	0	25	0	33.3	0	20	20
<i>Sargassum polycystum</i>	0	0	100	0	50	0	50.0	0	40	0
<i>Ulva lactuca</i>	0	0	100	0	100	0	100.0	0	60	0
<i>Nannochloropsis salina</i>	0	0	0	0	0	0	27.7	0	0	0
<i>Isochrysis galbana</i>	0	0	0	100	0	0	27.7	11.5	0	0
No plants (control)	0	0	0	0	0	0	16.6	0	0	0

the control (0 mg/l). *Gracilaria corticata* registered marginal decrease in GPP at 5 mg/l (9.0% less) and 10 mg/l (21.5% less) and an increase of 45% at 15 mg/l as well as 25 mg/l of ambient CO₂ levels over the control. However, *Ulva lactuca* did not show any marked difference. The planktonic algae registered higher GPP rates at 10 mg/l of CO₂ (111%) and at 15 mg/l CO₂ (84%). *Isochrysis galbana* showed 49% higher GPP only at 10 mg/l level and beyond this level registered considerable reduction in GPP (71% less at 15 mg/l and 89% less at 25 mg/l of CO₂) besides showing 84% less GPP at 5 mg/l concentration of CO₂ over the control.

Table 2. GPP (% over the control) of marine algae in different levels of dissolved CO₂

Species	CO ₂ level (mg/l)				
	0	5	10	15	25
<i>Gracilaria corticata</i>	100	91	78.5	145.2	146
<i>Sargassum polycystum</i>	100	224	0	252	309
<i>Ulva lactuca</i>	100	93	100.0	95	109
<i>Nannochloropsis salina</i>	100	89	211	184	89
<i>Isochrysis galbana</i>	100	16	149	29	11

The values presented in Table 1 were used to compute the CO₂ sequestering efficiency of the green, brown and red seaweeds along the Indian coasts as the standing stock of Indian seaweeds and their composition are known (Kaladharan and Reeta Jayasankar, 2003). The green seaweeds represented by *Ulva lactuca* are more efficient in carbon utilization than their brown and red counterparts (Table 3). The CO₂ emission rates of *Ulva* and *Sargassum* were observed as negligible and the net carbon utilisation of the Indian seaweed biomass is projected as 8687 t CO₂/ day.

Discussion

Although seaweeds and phytoplankton are considered as excellent sequestering agents of GHGs,

quantitative estimates were not available for the Indian seaweed biomass. Two culture systems are in use for CO₂ sequestration with planktonic algae—one is the open pond system and the other is the closed photobioreactor system and there is ongoing discussion regarding the merit of these two systems (Benemann, 1993; Pedroni *et al.*, 2001). We used closed photobioreactor system employing airtight, thick and clear polythene bags. The results obtained by us, though preliminary, indicate the capacity of the marine algae to utilize the excess CO₂ dissolved in the ambient water at levels 5 mg/l higher than the *in situ* levels in general and *Ulva lactuca* in particular, the commonly occurring seaweed along the intertidal zones of Indian coast. The emission of CO₂ through their respiratory activity is also minimum (0 to 20%) as these algae are capable of reutilizing the respiratory release of CO₂ within their cellular interspace for subsequent photosynthesis (Kanwisher, 1966).

The merit in the present investigation is that both the planktonic algae and macroalgae employed were representing different taxa such as green, brown and red algae. The methodology used was very simple but effective in the sense that within the same experimental set up CO₂ emitted or utilized and primary productivity could be monitored simultaneously by drawing samples for estimation through a hypodermic syringe where chances of contamination of extraneous air is minimum. This study also provides ample chance for quantifying the carbon sequestering efficiency of seaweeds as well as phytoplankton species.

Atmospheric CO₂ levels play a major role in the rate of carbon assimilation in terrestrial plants, and lower concentration of CO₂ promote higher photosynthesis (Forrester *et al.*, 1966). On the contrary, their marine counterparts, especially the

Table 3. Total estimated CO₂ absorbed (t/day) and emitted (t/day) by seaweed biomass along the Indian coasts

Type of seaweeds	Standing crop(t)*	Efficiency to absorb (mg/g/h)	CO ₂ absorbed (t/day)	Efficiency emit to (mg/g/h)	CO ₂ emitted (t/day)
Red algae	36523	1.60	584	1.0	365
Brown algae	41740	2.35	981	0	0
Green algae	182613	4.10	7487	0	0
Total	260876	8.05	9052	1.0	365

* Data from Kaladharan and Reeta Jayashanker (2003)

seaweeds registered higher rate of GPP (carbon assimilation) in increased levels of ambient CO₂ (Table 2) indicating the possibility of their unaltered carbon sequestering efficiency even in higher levels of dissolved CO₂.

It is estimated by Kaladharan and Reeta Jayasankar (2003) that the standing crop of seaweeds in the Indian waters was 2,60,876 tonnes, comprising 14% agar and carrageenan yielding seaweeds (Rhodophyceae), 16% algin yielding seaweeds (Phaeophyceae) and 70% green algae (Chlorophyceae). Taking these estimates into consideration, the CO₂ assimilation rate per day was computed for green algae represented by *Ulva lactuca*, brown algae represented by *Sargassum polycystum* and red algae represented by *Gracilaria corticata* (Table 3). As the green seaweeds are dominant, their carbon assimilation efficiency is high (Table 1). They are comparatively under-exploited as well as distributed throughout the intertidal zones to the euphotic zones of the neritic waters, and hence, sequester huge quantities of dissolved CO₂ from the coastal waters. Large scale mariculture of seaweeds especially of commercially important species can marginally reduce CO₂ concentration in the atmosphere as well as provide large harvestable biomass of raw material for phycocolloid industry.

Acknowledgements

The authors are grateful to the Director, CMFRI for constant encouragement and for providing necessary facilities. Field and laboratory assistance rendered by Sri Ch. Moshe is appreciated.

References

- Benemann, J. R. 1993. Utilization of carbon dioxide from fossil fuel- burning power plants with biological system. *Energy Conversion and Management*, 34 (9/11): 999-1004.
- Dye, J. F. 1958. Correlation of the two principal methods of calculating the three kinds of alkalinity. *J. Amer. Water Works Assoc.*, 50: 812-814.
- Forrester, M. L., G. Krotkov and C. D. Nelson. 1966. Effect of oxygen on photosynthesis, photorespiration and respiration in detached leaves 1. Soybean. *Plant Physiol.*, 41: 422-427.
- Gaarder, T. and H. M. Gran. 1927. *Rapp P.V. Reun. Comm. Int. Explor. Mer. Mediter.*, 42: 3-7.
- IPCC. 2007. Synthesis Report. Contribution of Working Groups of the Fourth Assessment Report of the Inter governmental Panel on Climate Change, WMD and UNEP, 397 pp.
- Kaladharan, P. and Reeta Jayasankar. 2003. Seaweeds. In: M. Mohan Joseph and A. A. Jayaprakash (Eds.) *Status of exploited marine fishery resources of India, Cent. Mar. Fish. Res. Inst., Cochin*, p. 228-239.
- Kanwisher, J. W. 1966. Photosynthesis and respiration in some seaweeds. In: Barnes, H. and George Allen (Eds.) *Some Contemporary Studies in Marine Science*, Unwin publishers, London, p. 407-420.
- Pedroni, P., J. Davison, H. Beckert, P. Bergmann and J. Benemann. 2001. A proposal to establish an international network on biofixation of CO₂ and greenhouse gas abatement with microalgae. *J. Ener. Environ. Res.*, 1(1): 136-150.
- Westlake, D. F. 1963. Comparisons of plant productivity. *Biol. Review*, 38: 385-425.
- Zou, D. 2005. Effects of elevated atmospheric CO₂ on growth, photosynthesis and nitrogen metabolism in the economic brown seaweed, *Hizikia fusiforme* (Sargassaceae, Phaeophyta). *Aquaculture*, 250: 726-735.

Received : 19/12/08
Accepted : 16/02/09