

# CMFRI bulletin 41



DECEMBER 1987

## **SEAWEED RESEARCH AND UTILIZATION IN INDIA**

**CENTRAL MARINE FISHERIES RESEARCH INSTITUTE**  
(Indian Council of Agricultural Research)  
P. B. No. 2704, E. R. G. Road, Cochin 682 031, India

## COMMON SEAWEED PRODUCTS

V. S. K. CHENNUBHOTLA, N. KALIAPERUMAL AND  
S. KALIMUTHU

The seaweeds are the only source for agar and algin. They are also used as food material, livestock feed and fertilizer in many parts of the world. The various products obtained from Indian seaweeds and their uses are dealt with here.

*Agar-Agar*

Agar-agar is a gelatinous substance obtained from the red algae like *Gelidium*, *Gelidiella* and *Gracilaria* and has a great commercial value. Agar is a colloidal carbohydrate present in the cell walls of some red algae and is a mixture of two polysaccharides, agarose and agaropection. Humm (1951) and Yaphe (1959) have defined agar as a gel-forming substance soluble in hot water and requiring one percent solution to set as a gel on cooling.

The important and commonly occurring agarophytes of India are *Gelidiella acerosa*, *Gracilaria edulis*, *G. crassa*, *G. verrucosa*, *G. corticata* and *G. foliifera*. The yield and physical properties of agar extracted from these red seaweeds are given in the Appendix. Among these seaweeds only *Gelidiella acerosa*, *Gracilaria edulis* and *G. crassa* are used at present as raw material for the production of agar-agar in India, since the yield and quality of agar are good and the plants are also available in harvestable quantities.

*Agaroids*

The gel-like extracts produced from certain types of red seaweeds are commonly known as agaroids. Carrageenan obtained from *Chondrus*, *Gigartina* and *Eucheuma* species come

under this group. The organic sulphate content is very much higher in these compounds and the chemical nature and properties of agaroids vary from that of agar. Pure solutions of agaroids are viscous and do not form gel when cooled. However, various inorganic and organic solutes alter the properties and increase the gelling power of agaroids.

Carrageenan-yielding plants have not been reported from Indian waters except for a rare and less abundant species, *Gigartina acicularis*, occurring in the intertidal region. But *Hypnea musciformis*, other species of *Hypnea*, *Spyridia*, *Sarconema*, *Acanthophora*, *Laurencia* and *Chondria* growing along the Indian coast give gel-like extracts.

*Algin or Alginic Acid*

Algin is the main polysaccharide occurring in the cell walls of brown algae. It consists of D-mannuronic acid and 2-guluronic acid in various proportions. The sodium, potassium and magnesium salts of alginic acid are soluble in water and they give viscous solutions without gel formation. Calcium alginate and other salts of copper, cobalt, mercury, etc are insoluble in water.

Species of *Sargassum*, *Turbinaria*, *Dictyota*, *Padina*, *Cystoseira*, *Hormophysa*, *Colpomenia*, *Spatoglossum* and *Stoechospermum* are some of the algin-yielding seaweeds occurring in Indian waters. The alginic acid content of these seaweeds is given in Table 10. Of these, *Sargassum* and *Turbinaria* are utilised as raw material for the manufacture of algin in India, since they are high-yielding varieties and also available in large quantities.

### Uses of Agar and Algin

Agar and algin are used in food, confectionary and dairy industries as gelling, stabilising and thickening agents, mainly in the manufacture of sweets, jellies, ice-creams, sherbats etc. They are also useful in a number of other industries.

**Agar :** Agar is extensively used in the making of food and medicines. The best known use of agar is as a solidifying agent in media used in bacteriological culture. It is also used as a stiffening agent in a number of food products, as a sizing material, and mucilage and in clarifying liquors. With its quality of keeping substances in suspension it goes in the manufacture of various pharmaceutical preparations, photographic film coatings and paints. It is employed in canning meat and in poultry, in laxative preparations, as a constituent of medical pills and capsules, in numerous pharmaceutical and cosmetic creams and jellies, as a dental-impression mould and as a lubricant for drawing tungsten in electrical bulbs.

**Algin :** Algin is also equally and extensively used in the preparation of various pharmaceutical, food and rubber products (natural and synthetic latex creaming and thickening, finished articles, automobile carpetting, electrical insulations, foam cushions, and rubber coating on tyres), textile products (size compound for cotton and rayon, textile print pastes and plastic laundry starch), adhesives (for all boards, paper bags, shipping containers, gummed tapes), paper products, food packages, pharmaceutical and detergent, packages, milk containers, butter cartons, frozen food packages, insulation boards, food wrappers, greaseproof paper and acoustical tiles) and miscellaneous products (paints, ceramic glazes, porcelain wares, leather finishers, autopolishes, welding-rod coatings, boiler compounds, batteryplate separators, wall-board-joint cement, beet-sugar processing and wax emulsions).

Some of the food products requiring agar and their method of preparation (Thivy, 1958) are given in Table 1.

Table 1  
*Food products requiring agar*  
(From Thivy, 1958)

Food stuff	Quantity of agar used	Method of addition
Ice-cream	1/8 teaspoonful (¼ g) per cup of ice-cream mix.	Dissolved in boiling water and added to warm ice-cream mix (Prevent it from melting soon)
Tomato sauce	½ teaspoonful (1 g) per 1b. of tomato sauce	Dissolved in boiling water and added to the sauce towards the end. boiling after adding agar should be avoided.
Jams, jelly, Marmalade	One level teaspoonful (2 g) per 1b. of these	Dissolved in boiling water and added to the sauce towards the end. Boiling after adding agar should be avoided.
Blancmange (without corn flour)	1½ level teaspoonful (3 g) per cup of milk with sugar	Dissolve agar in a small amount of water in a double boiler and pour in to warm milk, not vice versa.
Lime jelly	1½ level teaspoonful (3 g) per cup of water with sugar and lime juice	Dissolve agar in the water in a double boiler, add sugar and strain; keep aside and then, when somewhat cool, add lime-juice and pour into mould.

## Algal Proteins

Some green and red seaweeds such as *Ulva fasciata*, *U. rigida*, *Porphyra vietnamensis* and *Centroceras clavulatum* contain very rich proteins. These algal proteins have many essential amino acids including iodine-containing amino acids. Studies revealed that these seaweeds contain 16-30% of protein on dry weight basis and this amount is somewhat higher than that in other food materials such as cereals, eggs and fish (Visweswara Rao, 1964). Protein can be extracted from these seaweeds and as such dry powders of *Ulva*, *Porphyra*, *Acanthophora* etc. can be added to various foods deficient in protein or taken along with other food stuffs in small quantities.

## Seaweed as Food

Fresh, dried and processed seaweeds are utilised for human consumption. The algal carbohydrates are not easily digestible and the food value of the seaweeds depends on the minerals, trace elements, proteins and vitamins present in them. Many seaweeds such as species of *Caulerpa*, *Codium*, *Hydroclathrus*, *Sargassum*, *Porphyra*, *Gracilaria*, *Acanthophora* and *Laurencia* are used as food in Japan, Indonesia, China, Philippines and other countries of Indo-Pacific regions (Subba Rao, 1965; Levring *et. al.*, 1969; Michanek, 1975 and Chapman and Chapman, 1980). They are eaten as salad, curry, soup or vegetables. There are large industries in Japan using edible seaweeds like *Porphyra*. Thin algal sheets are prepared by washing and drying *Porphyra* plants and this forms an important food item in Japan.

Some of the edible seaweeds occurring in different localities along the Indian coast are species of *Ulva*, *Enteromorpha*, *Chaetomorpha*, *Caulerpa*, *Codium*, *Dictyota*, *Padina*, *Colpomenia*, *Hydroclathrus*, *Rosenvingea*, *Chnoospora*, *Sargassum*, *Turbinaria*, *Porphyra*, *Halymenia*, *Grateloupia*, *Gracilaria*, *Hypnea*, *Rhodymeina*, *Centroceras*, *Acanthophora*, and *Laurencia*. The important edible red seaweed *Porphyra* has been reported from Madras (Boergesen, 1937 b), Visakhapatnam and Cape Comorin (Uma-

maheswara Rao and Sreeramulu, 1963 and Umamaheswara Rao, 1973) and Goa Coast (Dhargaikar *et. al.*, (1981). The methods of preparing different recipes from seaweeds are given in detail by Chennubhotla *et. al.* (1981). The seaweed *Gracilaria edulis* is being used since decades for making gruel in the coastal areas of Tamil Nadu.

## Seaweed Meal

Seaweeds are cheap sources of minerals and trace elements. Hence the meals prepared from seaweeds can be given as supplements to the daily rations of the cattle, poultry and other farm animals. Seaweed meal can be obtained by grinding cleaned and washed seaweeds such as *Ulva*, *Enteromorpha*, *Sargassum*, *Padina*, *Dictyota*, *Gracilaria* and *Hypnea*. Thivy (1960) has described a simple method for the preparation of seaweed meal from *Gracilaria edulis* (*-G. lichenoides*). Seaweed meal can also be mixed with fish meal and used as a poultry feed. Seaweeds have been utilised as animal feed in some countries. Dave *et. al.* (1977) assessed the possibility of seaweeds being used as supplementary animal feed and they reviewed the feeding trials of animals with seaweeds conducted in Japan, Germany, U. K., Norway and other countries. The seaweed meal prepared from *Sargassum* and the results of its feeding trials on chicks, sheep and cattle are given by Dave *et. al.* (1979). Studies on feeding *Gracilaria* meal to white leghorn egg laying birds were made by Chaturvedi *et. al.* (1979), to find out the effect of algal-feed on the contents of egg. The results of this feeding experiment indicated that there was no significant difference in the number of eggs produced, total egg mass, internal quality of eggs and the body weight of the birds in the conventional ration group and those kept on ration with 5 and 10% *Gracilaria* meal. The *Gracilaria* meal at the level of 10% can be included in the ration of egg-laying birds, replacing yellow maize. Feeding trials replacing ragi (*Eleusine coracana*) with 0, 5, 10 and 15% of seaweed were conducted in unsexed day-old white leghorn chicks (Jagannathan and Venkatakrisnan, 1979) using six species of seaweeds commonly available in Tamil Nadu coast. 120 to 240

numbers of chicks were randomly allotted to the four diets with three to six replicates in each treatment. The trails were run for ten weeks. It is found that all these six seaweeds, particularly *Hypnea musciformis* and *Gracilaria edulis*, can be beneficially used to replace ragi at 5% level.

#### *Seaweed Manure*

Use of seaweeds as manure is a common practice in coastal areas throughout the world. In India it is used for coconut plantations especially in coastal Tamil Nadu and Kerala. The high amount of water soluble potash, other mineral and trace elements present in seaweeds are readily absorbed by plants and they control deficiency diseases. The carbohydrates and other organic matter present in seaweeds alter the nature of the soil and improve its moisture retaining capacity. Hence, large quantities of seaweeds including seagrasses such as *Cymodocea*, *Diplanthera*, *Enhalus* and *Halophila* can be used as manure in all parts of the country either directly or in the form of compost. A method for composting the seaweeds with cowdung has been described by Thivy (1958 and 1960). In the field trials conducted at the Central Marine Fisheries Research Institute, *Hypnea* compost when applied to bhendi, induced 73% increase in yield compared to cowdung and wood ash. Good results were also achieved with brinjal, tapioca, clustered beans, beans gouards, Amaranthus, Viridis lime, papaya and drumstick when manured with seaweed compost. Crotons and zinnias also grew well with seaweed treatment (Thivy, 1960). The nitrifiability of organic nitrogen from *Ulva lactuca* and drift seaweeds from Veraval was studied and found to be high compared to farmyard manure or a few other organic manures (Mehta *et. al.*, 1957). Application of seaweed manure can maintain a high level of available nitrogen in soil. The easy decomposability of seaweed organic matter is beneficial for the growth of soil micro organisms. According to these authors, seaweeds, especially the drift seaweed, which is a mixture of a variety of species cast ashore, can be a promising supplementary organic manure. The results of seaweed manurial trials on *Pennisetum typhoi-*

*des* (pearl millet) and *Arachis hypogea* (ground nut) are reported by Bokil *et. al.* (1972). The application of seaweed manure along with inorganic fertilizers have improved the quality of the produce. The use of seaweed manure had no significant influence over the yield of *Pennisetum typhoides* But the quality of its grains and fodder was favourably influenced. In the experiments on *Arachis hypogea* seaweed manure was comparable to that of farm yard manure and in general the performance of treatments in which seaweed manure was included was better than other treatments. With a view to find out the effect of seaweed manure on the uptake of inorganic nutrients by the wheat plant, a pot culture experiment was conducted (Bokil *et. al.*, 1974). The performance of the seaweed manure was found to be superior to the conventional manures, the performance of the seaweed manure is significantly better than that of farm yard manure due to the easy decomposibility of its carbonaceous matter and presence of micro-nutrients. The performance of brown seaweed manure, which contains higher proportion of alginic acid and analogous compounds, is relatively superior to that of drift seaweed manure with respect to both the yield and quality attributed. Regarding the quality of grains, the use of brown seaweed manure in conjunction with the inorganic fertilizers is significantly better than the others. Bhosle *et. al.* (1975) studied the seaweed extract on the growth of *Phaseolus vulgaris*. Marine algal extracts obtained from *Spatoglossum asperum*, *Ulva fasciata* and *Enteromorpha intestinalis* were found to promote germination in seeds and growth of seedlings of gram, ground nut and maize (Bukhari and Untawale, 1978). Dilute extracts were found to be more effective than the concentrated extracts. Foliar spray of *Spatoglossum* extract caused in an increase in the leaf size and better growth in *Hydrangia* sp. The method of preparation and properties of liquid seaweed fertilizer from *Sargassum* was given by Sreenivasa Rao *et. al.* (1979 a).

#### *Seaweed as a Source of Energy*

Sreenivasa Rao *et. al.* (1979 b) have conducted experiments on production of fuel gas for domestic use, utilizing *Sargassum* as raw

material. Digester design and operating parameters are given. According to them a mixture of about six micro-organisms mostly derived from marine environments were used in digesters. Addition of indole acetic acid stimulated anaerobic digesters. Salinity of the liquid above 20% was stated to be detrimental to production of fuel gas.

#### *Medicinal Uses of Seaweeds*

Seaweeds were considered to be of medicinal value in the Orient as early as 3000 B. C. The Chinese and Japanese used them in the treatment of goitre and other glandular diseases. Although the Romans believed seaweeds to be useless, they also used them to heal wounds, burns, scurvy and rashes. The British used *Porphyra* to prevent scurvy during long voyages.

Various red algae (particularly *Coralline officinalis*, *C. rubens* and *Alsidium heliminthocorton*) were employed as vermifuges in ancient times. Dulse is reported to be a laxative and also used to reduce fever. Several red algae (including *Chondrus crispus*, *Gracilaria*, *Gelidium* and *Pterocladia*) have been used to treat various stomach and intestinal disorders. The algae apparently absorb enough water and its water content helps relieving constipation and other associated discomforts. The stipes of *Laminaria cloustoni* have been used to aid in child birth by distending the uterus during labour. A number of species of marine algae have been found to have anticoagulant and antibiotic properties. Carrageenan may be useful in ulcer therapy and the alginates are found to prolong the "rate of activity" of certain drugs (Mathieson, 1969). Species of *Sargassum* were used for cooling and blood cleaning effect. *Hypnea musciformis* was employed as vermifuge or worm expelling agent and *Centroceras clavulatum* as cathartic agent. The iodine rich seaweeds such as *Asparagopsis taxiformis* and *Sarconema* can be used for controlling goitre disease caused by the enlargement of thyroid gland (Umamaheswara Rao, 1970).

Though the importance of different

seaweed products in pharmacology is known, the development of antimicrobial, antifungal and antiviral substances from seaweeds is still in an initial stage of research and development. Extracts from *Chondrus crispus* and *Gelidium cartilagineum* have been found to be active against influenza B and mumps virus, (Garber *et. al.*, 1958). Henriquez *et al.* (1979) assayed 33 species of Chilean marine algae for their antibacterial activity against *Sarcina lutea* ATCC 001, *Staphylococcus aureus* ATCC 65388 and *Bacillus subtilis* ATCC 6633. Some degree of antibacterial activity was found to be present in 17 of these 33 extracts. Caccamese *et. al.* (1980) tested the lipid extracts of more than 20 algae from eastern Sicily for antimicrobial activity against tobacco mosaic virus. Some of them mainly belonging to Dictyotales were found to be active. Caccamese *et. al.* (1981) tested lipid extracts of 13 algae from Eastern Sicily for antimicrobial activity against *Bacillus subtilis* and *Phoma tracheiphila* and for antiviral activity against tobacco mosaic virus. *Zanardine prototypus* and *Cystoseira balearica* exhibited the best antimicrobial and antiviral activity among the species tested. Blunden *et. al.* (1981) examined the extracts of British marine algae for anti-influenza virus activity based on inhibition of influenza neuraminidase. The antibacterial and antifungal activity of Indian seaweed extracts (Sreenivasa Rao *et. al.*, 1979 c; Sresnivasa Rao and Shelat, 1979 and Sreenivasa Rao and Parekh, 1981) and also the effect of seaweed extracts on *Mycrobacterium tuberculosis* (Sreenivasa Rao *et. al.*, 1979 d) have been studied. Antibiotic substance isolated from *Enteromorpha* effected complete inhibition of growth of the tubercle bacilli in the cultures. Naqvi *et. al.*, (1981) examined the extracts of 25 seaweeds from Indian coast for antiviral, antibacterial, antifungal, antiprotozoal, antifertility activities and a wide range of pharmacological activities. Significant biological activity was obtained in 13 species of seaweeds, the most promising activity being 100% antifertility (anti-implantation) activity observed in 3 species namely *Padina tetrastratica*, *Gelidiella acerosa* and *Acanthophora spicifera*.

BIBLIOGRAPHY

- AGADI, V. V. AND A. G. UNTAWALE. 1978. Marine algal flora of Goa coast. *Seaweed Res. Util.*, 3 (1 & 2): 56-70.
- AGADI, V. V., N. B. BHOSLE AND A. G. UNTAWALE. 1978. Metal concentration in some seaweeds of Goa (India). *Bot. Mar.*, 21 (4): 247-250.
- AGHARKAR, S. P. 1923. The present position of our knowledge of the aquatic flora of India. *Jour. Indian. bot. Soc.*, 3: 252-260.
- ANON, 1983. Proven Technology 7. Technology of cultured seaweed production. *Mar. Fish. Infor. Ser. T & E. 54*: 19-20.
- BACKGROUND, A 1976. "Nori" farming in Japan (cultivation of the edible seaweeds of the genus *Porphyra*). In: *Farming marine organisms low in the food chain. Development in Aquaculture and Fisheries Science*. Vol. 1 (Ed. P. Korringa) Elsevier Scientific Publishing Co. Amsterdam. pp 17-48.
- BALAKRISHNAN Nair, N., N. SHOBA AND M. ARUNACHALAM. 1982. Algae from southern Kerala coast. *Indian J. Mar Sci.*, 11 (3): 266-269.
- BARDACH, J. E. J., H. RYHTER AND W. O. MCLARNEY. 1972. Seaweed culture. In: *Aquaculture. The Farming and husbandry of Freshwater and Marine organisms*. Wiley-Interscience. A division of John Wiley & Sons, Inc. New York pp 790-840.
- BHANDERI, P. P. 1974 a. An estimate of the iodine yielding seaweed *Asparagopsis taxiformis* (Delile) Collins and Harvey from some subtidal reefs of Saurashtra coast. *J. mar. biol. Ass. India.*, 16 (1):288-289.
- BHANDERI, P. P. 1974 b. Culture of the agar yielding seaweeds on ropes from Gujarat. *J. mar. biol. Ass. India*, 16 (3):847-849.
- BHANDERI, P. P. AND Y. B. RAVAL. 1975. Possibility of seaweed cultivation along the Gujarat coast. *Seafood Export Jour.*, 7 (12):33-36.
- BHANDERI, P. P. AND Y. A. TRIVEDI. 1975. Seaweed resources of Hanumandandi reef and Vumani reef near Okha Port. *Indian J. mar. Sci.*, 4 (1):97-99.
- BHANDERI, P. P. AND Y. A. TRIVEDI. 1977. Rope culture of algin yielding seaweed *Hormophysa triquetra* (Linnaeus) Kuetzing. *Bot. Mar.* 20 (3):183-185.
- BHOSLE, N. B., V. K. DHARGALKAR AND A. G. UNTAWALE. 1975. Effect of seaweed extract on the growth of *Phaseolus vulgaris* L. *Indian J. mar. Sci.*, 4:207-210.
- BISWAS, K. 1932. Census of Indian algae. Scope of algological studies in India. *Rev. Algol.*, 6:197-219.

- BISWAS, K. 1934. Progress of algological studies in India. *Curr. Sci.*, 3:237-241.
- BISWAS, K. 1945. A general review of the marine algae of the western coast of India. *Jour. Bombay nat. Hist. Soc.* 45:515-530.
- BLACK, W. A. P. 1954. Constituents of the marine algae. *Ann. Rep. Chem. Soc.*, 50:322-335.
- BLANCO, G. J. 1973. Status and problems of coastal aquaculture in the Philippines. In: *Coastal Aquaculture in the Indo-Pacific Region* (Ed. T. V. R. Pillay) F. A. O. Fishing News (Books) Ltd., pp. 60-67.
- BLUNDEN, G., C. J. BARWELL, K. J. FIDGEN AND K. JEWERS. 1981. A survey of some British marine algae for anti-influenza virus activity. *Bot. Mar.*, 14 (5) 267-272.
- BOERGESEN, F. 1930. Some Indian green and brown algae especially from the shores of the Presidency of Bombay. *Jour. Indian bot. Soc.*, 9:151-174.
- BOERGESEN, F. 1931. Some Indian Rhodophyceae especially from the shores of the Presidency of Bombay—I. *Kew. Bull.*, No. 1: 1-24.
- BOERGESEN, F. 1932a. Some Indian Rhodophyceae especially from the Presidency of Bombay—II. *Kew. Bull.*, No. 3: 113-134.
- BOERGESEN, F. 1932 b. Some Indian green and brown algae especially from the Presidency of Bombay. *Jour. Indian. bot. Soc.*, 11: 51-70.
- BOERGESEN, F. 1933 a. Some Indian Rhodophyceae especially from the Presidency of Bombay III. *Kew. Bull.*, No. 3: 113-142.
- BOERGESEN, F. 1933 b. Some Indian green and brown algae from the Presidency of Bombay. *Jour. Indian bot. Soc.*, 12: 1-16.
- BOERGESEN, F. 1934 a. Some Indian Rhodophyceae especially from the Presidency of Bombay-IV. *Kew. Bull.*, No. 4: 1-30.
- BOERGESEN, F. 1934 b. Some marine algae from the northern part of the Arabian Sea with remarks on their geographical distribution. *Kgl. Dansk. Vidensk. Selskab Biol. Meddel.* 11 (6): 1-72.
- BOERGESEN, F. 1935. A list of marine algae from Bombay. *Kgl. Dansk. Vidensk. Selskab Biol Meddel.*, 12 (2): 1-64.
- BOERGESEN, F. 1937 a. Contributions to a South Indian Marine Algal flora-1. *Jour. Indian bot. Soc.*, 16: 1-56.
- BOERGESEN, F. 1937 b. Contributions to a South Indian Marine Algal flora-II. *Jour. Indian bot. Soc.*, 16: 311-357.
- BOERGESEN, F. 1938. Contributions to a South Indian Marine Algal flora-III. *Jour. Indian bot. Soc.*, 17: 205-242.
- BOKIL, K. K., V. C. MEHTA AND D. S. DATAR. 1972. Seaweed as manure. III, Field manurial trials on *Pennisetum typhoides* S. H. (Pearl Millet) and *Arachis hypogaea* (Groundnut). *Bot. Mar.*, 15 (3): 148-150.



- BOKIL, K. K., V. C. MEHTA AND D. S. DATAR. 1974. Seaweed as manure: II. Pot culture manurial experiments on wheat. *Phykos*, 13 (1): 1-5.
- BOSE, J. L., KARIMULLAH AND S. SIDDIQUE. 1943. Manufacture of agar in India. *J. Sci. Industr. Res.*, (India) 1: 98.
- BURELLY, P. 1968. Les algues d'eau douce Bou bee et. Cie, Paris. Tomes 1, 2 and 3.
- BUKHARI, S. S. AND A. G. UNTAWALE. 1978. Seaweeds as liquid fertilizer and foliar spray. *Seaweed Res. Util.*, 3 (1&2): 71-78.
- CACCAMESE, S., R. AZZOLINA, G. FURNARI, M. CORMACI AND S. GRASSO. 1980. Antimicrobial and antiviral activities of extracts from Mediterranean algae. *Bot Mar*, 23 (5): 285-288.
- CACCAMESE, S., R. AZZOLINE, G. FURNARI, M. CORMACI AND S. GRASSO. 1981. Antimicrobial and antiviral activities of some marine algae from eastern Sicily. *Bot Ma.*, 24 (7): 365-367.
- CAMERON, M. C., A. G. ROSS AND E. G. V. PERCIVAL. 1948. Methods of the routine estimation of mannitol, alginic acid and combined fucose in seaweeds. *Jour. Soc. Chem. Ind.* London, 67: 161-164.      ◆
- CHACKO, P. I. AND C. MALU PILLAI. 1958. Studies on utilisation of the seaweed resources of Madras State. *Contr. Mar. Biol. St. Krusadai Island*, 6: 1-12.
- CHACKO, P. I., S. MAHADEVAN AND R. GANESAN, 1955. A guide to the field study of the fauna and flora of Krusadai Island, Gulf of Mannar. *Contr. Mar. Biol. St. Krusadai Island*, 3: 1-16.
- CHADEFAUD, M. 1960. *Traite de Botanique*, Tome 1, Masson et. cie, Paris.
- CHAKRABORTY, D. 1945. Agar-agar manufacture from *Gracilaria confervoides*. *Jour. Proc. Inst. Chem.* (India), 17: 188.
- CHAPMAN, V. J. 1962. *The algae*. London.
- CHAPMAN, V. J. AND D. J. CHAPMAN. 1975. *The algae*. English Language Book Society and Macmillan, 497 pp.
- CHAPMAN, V. J. AND D. J. CHAPMAN, 1980. *Seaweeds and Their Uses*. Third Edition, Chapman and Hall, London. pp 62-97.
- CHATURVEDI, D. K., S. S. VERMA AND S. P. KHARE. 1979. Studies on feeding marine algae (Gracilaria meal) to laying white leghorn birds. *Proc. Int. Symp. Marine Algae of the Indian Ocean Region*, CSMCRI, Bhavnagar, India. p. 51 (Abstract).
- CHAUHAN, V. D. 1970. Variation in alginic acid content with growth stages in two species of *Sargassum*. *Bot. Mar.*, 13 (1): 57-58.
- CHAUHAN, V. D. 1972. Physiological ecology of the early stages of *Sargassum swartzii* (Turner) C. Ag. *Bot. Mar.* 15 (1): 49-51.
- CHAUHAN, V. D. 1978. Report on the survey of marine algae resources of Maharashtra coast. *Salt. Res. Ind.*, 14 (1): 1-10.

- CHAUHAN, V. D. AND V. KRISHNAMURTHY. 1967. Observations on the output of oospores, their liberation, viability and germination in *Sargassum swartzii* (Turn) C. Ag. *Proc. Semi. Sea Salt and Plants*, CSMCRI, Bhavnagar, pp. 197-201.
- CHAUHAN, V. D. AND V. KRISHNAMURTHY. 1968. An estimate of algin bearing seaweeds in the Gulf of Kutch. *Curr. Sci* 37: 648.
- CHAUHAN, V. D. AND O. P. MAIRH. 1978. Report on the survey of marine algae resources of Saurashtra coast, India. *Salt. Res. Ind.*, 14 (2): 21-41.
- CHAUHAN, V. D. AND H. V. JOSHI. 1979. Effect of Indole-3-Acetic acid and Gibberellic acids on the early growth of *Sargassum*. *Proc. Int. Seaweed Symp. Marine Algae of the Indian Ocean Region*. CSMCRI, Bhavnagar, India. p. 23 (Abstract).
- CHEN, T. P. 1976. Culture of *Gracilaria*. In: *Aquaculture practices in Taiwan*. Page Bros. (Norwich) Ltd. pp. 145-149.
- CHENNUBHOTLA, V. S. K. 1976. Seaweed Culture. *Indian Science Congress 63rd Session, Waltair (Abstract)*.
- CHENNUBHOTLA, V. S. K. 1977. Food from the sea: food from the seaweeds. *Seafood Export Jour.*, 9 (3): 1-4.
- CHENNUBHOTLA, V. S. K., M. NAJMUDDIN AND BIDYADHAR NAYAK. 1977 a. A comparative study of the yield and physical properties of agar-agar from different blends of seaweeds. *Seaweed Res. Util.*, 2 (2): 87-90.
- CHENNUBHOTLA, V. S. K., S. KALIMUTHU, N. KALIAPERUMAL AND J. R. RAMALINGAM. 1977 b. Studies on the growth variation, alginic acid and mannitol contents in *Padina gymnospora* (Kuetzing) Vickers. *Seaweed Res. Util.* 2 (2): 91-94.
- CHENNUBHOTLA, V. S. K., S. KALIMUTHU, M. NAJMUDDIN AND M. SELVARAJ. 1977 c. Field culture of *Gelidiella acerosa* in the inshore waters of Gulf of Mannar. Supplement to *Jour. Phycol.* Vol. 13, Abstract No. 454.
- CHENNUBHOTLA, V. S. K., N. KALIAPERUMAL AND S. KALIMUTHU. 1978 a. Culture of *Gracilaria edulis* in the inshore waters of Gulf of Mannar (Mandapam). *Indian J. Fish.*, 25 (1 & 2): 228-229.
- CHENNUBHOTLA, V. S. K., N. KALIAPERUMAL AND S. KALIMUTHU. 1978 b. Seasonal changes in growth, fruiting cycle and oospore output in *Turbinaria conoides* (J. Ag.) Kuetzing. *Bot. Mar.*, 21 (1): 67-69
- CHENNUBHOTLA, V. S. K., KALIMUTHU, M. NAJMUDDIN, R. PANIGRAHY AND M. SELVARAJ. 1979. Seasonal variation in growth, yield of agar-agar and its physical properties in some agarophytes of Tamil Nadu Coast. *Proc. Int. Symp. Marine Algae of the Indian Ocean Region*. CSMCRI, Bhavnagar, India p. 41 (Abstract).
- CHENNUBHOTLA, V. S. K., N. KALIAPERUMAL AND S. KALIMUTHU. 1981. Seaweed recipes and other practical uses of seaweeds. *Seafood Export Jour.*, 13 (10): 9-16
- CHENNUBHOTLA, V. S. K., N. KALIAPERUMAL, S. KALIMUTHU, M. SELVARAJ, J. R. RAMALINGAM AND M. NAJMUDDIN. 1982. Seasonal changes in growth and alginic

- acid and mannitol contents in *Sargassum ilicifolium* (Turner) J. Agardh and *S. myriocystum* J. Agardh. *Indian J. Mar. Sci.*, 11 (2): 195-196.
- CHENNUBHOTLA, V. S. K. 1981. *Status of seaweed industry in India*. U N D P Report on the Training Course on *Gracilaria* algae of the South China Sea Fisheries Development and co-ordinating Programme, Manila, Philippines. No SCS/Gen/81/29: 139-145.
- CHENNUBHOTLA, V. S. K., N. KALIAPERUMAL, S. KALIMUTHU AND PVR. NAIR. 1983. Biology of the economically important Indian seaweeds. Collected Abstracts of the International Seaweed Symposium, Qingdao, China.
- CHENNUBHOTLA, V. S. K., S. KALIMUTHU, M. NAJMUDDIN, R. PANIGRAPHY AND M. SELVARAJ. 1986. Changes in growth and phycocolloid content of *Gelidium acerosa* and *Gracilaria edulis*. *Seaweed Res. Utiln.* 9 (1 & 2): 45-48.
- CHENNUBHOTLA, V. S. K., N. KALIAPERUMAL, J. R. RAMALINGAM AND S. KALIMUTHU. 1986. Growth, Reproduction and spore output in *Gracilaria foliifera* (Forsk.) Boergesen and *Gracilariaopsis sjoestedtii* (Kylin) Dawson around Mandapam. *Indian J. Fish.* 33 (1): 76-84.
- CHENNUBHOTLA, V. S. K., S. KALIMUTHU AND M. SELVARAJ. 1986. Seaweed culture—its feasibility and industrial utilization *Proc. Symp. Coastal Aquaculture* 4: 1206-1209.
- CHENNUBHOTLA, V. S. K., B. S. RAMACHANDRUDU, P. KALADHARAN AND S. K. DHARMARAJ. 1987. Seaweed resources of Kerala coast. Seminar on Fisheries Research and Development in Kerala. Trivandrum (Abstract).
- CHENNUBHOTLA, V. S. K., M. NAJMUDDIN, J. R. RAMALINGAM AND N. KALIAPERUMAL. 1987. Biochemical composition of some marine algae of Mandapam coast (South India). Symposium on Research and Development in Marine Fisheries. Mandapam camp.
- CHENNUBHOTLA, V. S. K., N. KALIAPERUMAL AND M. S. RAJAGOPALAN. 1987. Seaweed culture in India—An appraisal. *Ibid.*
- CHIDAMBARAM, K. AND M. M. UNNY. 1947. Note on the value of seaweeds as manure. *Madras Agri. Jour.* (July).
- CHIDAMBARAM, K. AND M. M. UNNY. 1953. Note on the value of seaweeds as manure. *Int. Seaweed Symp.*, pp. 67-68.
- CHRISTENSEN, T. 1962. *Alger*. In *systematisk Botanik* (Ed. T. W. Bocher, M. Lange and T. Sorenson) Vol. 2. No. 2. Munkagaard, Copenhagen. 178 pp.
- DAVE, H. M., V. SITAKARA RAO AND U. K. TIPNIS. 1969. Iodine content of marine algae from Saurashtra coast. *Phykos*, 8: 68-70.
- DAVE, M. J. AND R. G. PAREKH. 1975. Protein content of green seaweeds from Saurashtra coast. *Salt. Res. Indi.*, 11 (2): 41-44.
- DEVE, M. J., S. K. GARG AND E. R. R. IYENGAR. 1977. Assessment of the possibility of seaweeds to be utilised as supplementary animal feed. *Salt. Res. Ind.*, 13 (1&2): 33-40.

- DAVE, M. J., R. G. PAREKH, S. K. GARG AND D. J. METHA. 1979. Preparation of seaweed meal for the feeding of farm animals. *Salt Res. Ind.*, 15 (2): 34-38.
- DAVIDSON, F. E. 1950. The effect of auxin on the growth of marine algae. *Amer Jour. Bot.*, 37: 502-510.
- DAWSON, E. Y. 1966. *Marine Botany: An introduction*. Holt, Rinehart and Winston Inc, New York, 371 pp.
- DE JUSSIEU, A. L. 1789. *Genera plantarum secundum ordines naturales disposita*, Paris. 498 pp.
- DESAI, B. N. 1967. Seaweed resources and extraction of alginate and agar. *Proc. Semi Sea Salt and Plants*, CSMCRI, Bhavnagar, pp. 343-351.
- DEVEAU, L. E. AND J. R. CASTLE. 1979. The industrial development of farmed marine algae. The case history of *Euchenuma* in the Philippines and USA. In: *Advances in Aquaculture* (Ed. T. V. R. Pillay and Wm. A. Dill) F. A. O. Fishing News Books Ltd. England. pp 410-415.
- DHANDUKIA, M. M. AND R. SESHADRI. 1969. Arsenic content in marine algae. *Phykosa*, 8: 108-111.
- DHARGALKAR, V. K. 1979. Biochemical studies on *Ulva reticulata* Forsskal. *Proc. Int. Symp. Marine Algae of the Indian Ocean Region*, CSMCRI, Bhavnagar, p. 40 (Abstract)
- DHARGALKAR, V. K., T. G. JAGTAP AND A. G. UNTAWALE. 1980. Biochemical constituents of seaweeds along the Maharashtra coast. *Indian. J. Mar. Sci.*, 2 (4): 297-299.
- DHARGALKAR, V. K., V. V. AGADI AND A. G. UNTAWALE. 1981. Occurrence of *Porphyra vietnamensis* (Bangiales, Rhodophyta) along the Goa coast. *Mahasagar*, 14 (1): 75-77.
- DIXIT, S. C. 1964. Species list of Indian marine algae determined by Boergesen. *J. Uni. Bombay*, 32 (2-5): 1-23.
- DIXIT, S. C. 1968. Species list of Indian marine algae- II *J. Univ. Bombay*, 36 (3-5): 9-24.
- DODGE, J. D. 1969. A review of the fine structure of algal eye spots. *Br. phycol. J.* 4 199-210.
- DOSHI, Y. A. AND P. SREENIVASA RAO. 1967 a. Stable agar by gamma irradiation *Nature* 216: 931.
- DOSHI, Y. A. AND P. SREENIVASA RAO. 1967 b. Radiation induced enhancement of gel strength in red seaweeds. *Indian Jour Chem.*, 5: 342-343.
- DOSHI, Y. A., P. V. RAJU AND P. SREENIVASA RAO. 1968. A relation between the sulphate content in red seaweeds and the gel strength of agar. *Sci. Cul.*, 34: 493.
- DOUGHERTY, E. G. AND ALLEN, M. B. 1960. In: *Comparative biochemistry of photoreactive systems* (Ed. M. B. Allen). Academic Press, New York and London. pp. 129-143.
- DURAIRAJ, S., K. G. JOSEPH AND M. KINGSLEY LAINE. 1978. Preparation of sodium alginate with improved viscosity. *Seaweed Res. Util*, 3 (1&2): 5-8.

- EICHLER, A. W. 1886. Syllabus der Vorlesungen über speciella and Medicinisch-pharmaceutische Botanik, 4th Ed. Berlin
- ENDLICHER, S. 1836. Genera plantarum secundum ordines naturales disposita. Vindbonae
- FELDMANN, G. 1963. In *Precis de Botanique*. (Ed. M. Chadeaud and M. Emberger) Masson et. Cie, Paris. pp 83-249.
- FOTT, B. 1959. Algenkunde, Gustav Fisher, Jena.
- FRITSCH, F. E. 1935. *The structure and reproduction of algae*, Vol. I. University Press, Cambridge. pp. 1-791.
- FURUKAWA, A. 1973. Present status of Japanese marine aquaculture. In: *Coastal Aquaculture in Indo-Pacific Region* (Ed. T. V. R. Pillay). F. A. O Fishing News (Books) Ltd., pp. 29-47.
- GARBER, P., J. D. DUTCHER, E. G. ADAMS AND J. R. SHERMAN. 1958. Protective effects of seaweed extracts for chicken embryos infected with influenza B or mumps virus. *Proc. Soc. Exp. Biol. Med.*, 99. 590-593.
- GOLDSTEIN, M. E. 1973. Regeneration and vegetative propagation of the agarophyte *Gracilaria debilis* (Forsk.) Boergesen. (Rhodophyceae) *Bot. Mar.*, 26 (4): 226-228.
- GOPALAKRISHNAN, P. 1969. Some marine algae from the Gulf of Kutch. *Phykos*, 8: 61-67.
- GOPALAKRISHNAN, P. 1970. Some observation on the shore ecology of the Okha coast. *J. mar. biol. Ass. India*. 12 (1 & 2): 15-34.
- GUIST, G. G., C. J. DAWES AND J. R. CASTLE. 1982. Mariculture of the red seaweed, *Hypnea musciformis*. *Aquaculture*, 28 (3, 4): 375-384.
- HARVEY, W. H. 1836. Algae. In *Flora Hibernica*, Mackay, J. T. Dublin.
- HENRIQUEZ, P., A. CANDIA, R. NORAMBUENA, M. SILVA AND R. ZEMELMAN. 1979. Antibiotic properties of marine algae. II. Screening of Chilean marine algae for antimicrobial activity. *Bot. Mar.*, 22 (7): 461-463.
- HORNELL, J. 1918. *Report on the further development of fishery resources of Baroda State*.
- HUMM, H. J. 1951. The red algae of economic importance. Agar and related phycocolloids. In: *Marine Products of Commerce* (Ed. Tressler, D. K.). New York.
- IYENGAR, M. O. P. 1957. Algology in progress of science in India. Sn. VI. Botany. *Natn. Inst. Sci. India*. New Delhi: 229-251.
- JAGANNATHAN, V. AND R. VENKATAKRISHNAN. 1979. Nutritional investigations of seaweeds in chick ration. *Proc. Int. Symp on Marine Algae of the Indian Ocean Region CSMCRI, Bhavnagar, India* pp. 49-51 (Abstract).
- JAMES, P. S. B. R., V. S. K. CHENNUBHOTLA AND RODRIGO. 1980. Studies on the fauna associated with the cultured seaweed *Gracilaria edulis*. *Proc Symp. Coastal Aquaculture, M. B. A. I., Cochin, India*, p. 111 (Abstract)

- JOSEPH, I. AND S. MAHADEVAN, 1948. Production of agar-agar. *Dept. Res. Univ. Travancore, Rep. for Septen.*, pp. 55-60.
- JOSEPH, I., K. GANAPATHY AND S. RAMAMURTHY. 1948. Recoverable iodine from Indian *Sargassum*. *Dept. Res. Univ. Travancore, Rep. for Septen.*, pp. 60-61.
- JOSHI, A. C. 1949. Indian Botany; present position and prospects. Presidential Address. *Four. Indian bot. Soc.*, 28: 1-15.
- JOSHI, H. V. AND V. KRISHNAMURTHY. 1971. The species of *Enteromorpha* from India *Bot. J. Linn. Soc.*, 65 (1): 119-128.
- KALE, S. R. AND KRISHNAMURTHY. 1967. Effect of different media on the germlings of *Ulva lactuca* var. *rigida*. *Phykos*, 6 (1 & 2): 32-35.
- KALIAPERUMAL, N. AND M. UMAMAHESWARA RAO. 1975. Growth, fruiting cycle and oospore output in *Turbinaria decurrens* Bory. *Indian J. Fish.*, 22 (1 & 2): 225-230.
- KALIAPERUMAL, N. AND S. KALIMUTHU. 1976. Changes in growth, reproduction, alginic acid and mannitol contents in *Turbinaria decurrens* Bory. *Bot. Mar.*, 19: 157-159.
- KALIAPERUMAL, N. AND M. UMAMAHESWARA RAO. 1981. Studies on the standing crop and phycocolloid of *Gelidium pusillum* and *Pterocladia heteroplata* *Indian J. Bot.*, 4 (2): 91-95.
- KALIAPERUMAL, N. AND M. UMAMAHESWARA RAO. 1982. Seasonal growth and reproduction of *Gelidiopsis variabilis* (Greville) Schmitz *J. Exp. Mar. Biol. Ecol.*, 61: 265-270.
- KALIAPERUMAL, N., V. S. K. CHENNUBHOTLA AND S. KALIMUTHU. 1977. Growth, reproduction and liberation of oospores in *Turbinaria ornata* (Turner) J. Agardh. *Indian J. Mar. Sci.*, 6 (2): 178-179.
- KALIMUTHU, S. 1980. Variations in growth and mannitol and alginic acid contents of *Sargassum myriocystum* J. Agardh *Indian J. Fish.*, 27 (1 & 2): 265-266.
- KALIMUTHU, S., V. S. K. CHENNUBHOTLA, M. SELVARAJ, M. NAJMUDDIN AND PANIGRAHY, 1980. Alginic acid and mannitol contents in relation to growth in *Stoechospermum marginatum* (C. Agardh) Kuetzing. *Indian Fish*, 27 (1 & 2): 267-269.
- KANNAN, L. AND K. KRISHNAMURTHY. 1978. A survey of the algae of the Porto-Novo region (Coromandel Coast. Bay of Bengal). *Seaweed. Res. Util.* 3 (1 & 2): 1-4.
- KAPPANNA, A. N. AND V. SITAKARA RAO. 1932. Iodine content of marine algae from Gujarat coast. *Jour. Sci. Indust. Res. (India)*, 21: 559-560.
- KAPPANNA, A. N. AND A. VISWESWARA RAO. 1963. Preparation and properties of agar-agar from Indian seaweeds. *Indian Jour. Tech.* 1: 224.
- KAPPANNA, A. N., A. VISWESWARA RAO AND I. C. MODY. 1962. Alginic acid content of some of the brown seaweeds of Sourashtra coast. *Curr. Sci.*, 31: 463-464.
- KARUNAKAR, P. D., M. S. RAJU AND S. VARADARAJAN. 1948. Manufacture of agar-agar from seaweed, *Gracilaria lichenoides*. *Indian Vet. J.*, 24: 274.

- KOSHY, T. K. AND C. C. JOHN. 1948. Survey of *Gracilaria* resources of Travancore coast. *Dept. Res Unia. Travancore. Rep. for Septen..* pp 53-55.
- KOW, T. A., Y. S. LING AND T. W. HIN. 1973. Experiments in coastal aquaculture in Singapore. In: *Coastal Aquaculture in the Indo-Pacific Regions* (Ed. V. V. R. Pillay) F. A. O. Fishing News (Books) Ltd., England pp. 375-383.
- KRISHNAMURTHY, V. 1967. Marine algal cultivation—necessity, principles and problems. *Proc. Semi. Sea Salt and Plants* CSMCRI, Bhavnagar. pp. 327-333.
- KRISHNAMURTHY, V. 1980. The marine algae of Tiruchendur, South India. *Seaweed Res. Util.*, 4 (1): 49-58.
- KRISHNAMURTHY, V. AND H. V. JOSHI. 1969. The species of *Ulva* Indian waters. *Bot J. Linn. Soc.* 62: 123-130.
- KRISHNAMURTHY, V. AND H. V. JOSHI. 1970. A Check-list of Indian Marine Algae, CSMCRI, Bhavnagar. pp. 1-36.
- KRISHNAMURTHY, V. R. VENUGOPAL, J. G. THIAGARAJ AND H. N. SHAH. 1967. Estimating drift seaweeds on the Indian coasts. *Proc. Semi. Sea Salt and Plants*, CSMCRI, Bhavnagar, pp. 315-320.
- KRISHNAMURTHY, V., P. V. RAJU AND R. VENUGOPAL. 1969. An aberrant life history in *Gracilaria edulis* (Gmel.) Silva and *Gracilaria corticata* J. Ag. *Curr. Sci.*, 38 (14): 343-344.
- KRISHNAMURTHY, V., P.V. RAJU AND P. C. THOMAS. 1975. On augmenting seaweed resources of India. *J mar. biol. Ass. India.*, 17. (2): 181-185.
- KUROGI, M. 1963. Recent laver cultivation in Japan. *Fishing News Intl.*, 2 (3): 269-274.
- LANGALIA, J. K., K. SESHADRI AND D. S. DATAR. 1967. The alkali contents of the marine algae. *Proc. Semi Sea. Salt and Plants*, CSMCRI, Bhavnagar pp. 289-295.
- LEVRING, T., H. A. HOPPE AND O. J. SCHMID. 1969. *Marine Algae. A survey of Research and Utilization.* Gram, be Gruyter & Co., Hamburg, pp. 1-421.
- LEWIS, E. J. 1962 a. Studies on the proteins, peptides, and free aminoacid contents in some species of *Padina* from south-eastern coast of India. *Curr Sci.*, 31. 90-92.
- LEWIS, E. J 1962 b. Studies on the proteins peptides and free aminoacid contents in some species of brown algae from south-eastern coast of India. *Rev. Algal.*, 6: 209-216.
- LEWIS, E. J. 1963 a. Studies on the proteins, peptides and free aminoacid contents in some species of marine algae from south-eastern coast of India. *Rev., Algal.*, 7: 15-25.
- LEWIS, E. J. 1963 b. The proteins, peptides and free aminoacid composition in species of *Acanthophora* from south east coast of India *Rev., Algal.*, 7: 237-241.
- LEWIS, E. J. 1963 c. Studies on the proteins, peptides and free aminoacid contents in some species of red algae from south-eastern coast of India. *Proc. natn. Inst. Sci. India*, 29: 137-145.
- LEWIS, E. J. 1963 d. Studies on fortnightly analysis of the proteins, peptides and free aminoacids in some marine algae from Bombay. *Proc. natn. Inst. Sci. India*, 29: 363-286.

- LEWIS, E. J. 1967. A review of protein, peptide, and free aminoacid contents of Indian Marine Algae. *Proc. Semi. Sea Salt and Plants*, CSMCRI, Bhavnagar, pp. 296-308.
- LEWIS, E. J. AND E. A. GONZALVES. 1959 a. Studies on the free aminoacid contents of some marine algae from Bombay. *Jour. Univ. Bombay*, 28: 1-5.
- LEWIS, E. J. AND E. A. GONZALVES. 1959 b. Aminoacid contents of the erect and creeping fronds of species of *Caulerpa* from Bombay. *Jour. mar. biol. Ass. India*, 1: 54-56.
- LEWIS, E. J. AND E. A. GONZALVES. 1959 c. Studies on the free aminoacid contents of species of *Caulerpa* from Bombay. *Jour. mar biol Ass. India*, 1: 203-205.
- LEWIS, E. J. AND E. A. GONZALVES. 1960. Aminoacid contents of some marine algae from Bombay. *New Phytol*, 59: 109-115.
- LEWIS, E. J. AND E. A. GONZALVES 1962 a. Studies on the protein, peptide and free aminoacids in cystocarpic and tetrasporic plants of *Agardhiella roubusta* from Bombay. *New Phytol.*, 61: 288-290.
- LEWIS, E. J. AND E. A. GONZALVES. 1962 b. The protein, peptide and aminoacid contents of some species of marine algae from Bombay. *Ann. Bot. N. S.*, 26. 301-316.
- LEWIS, E. J. AND E. A. GONZALVES. 1962 c: Periodic studies of the proteins, peptides and free aminoacid in *Enteromorpha prolifera*, f. *capillaris* and *Ulva lactuca* var. *rigida*. *Ann. Bot. N. S.*, 26: 317-327.
- LING, S. W. 1973. A review of the status and problems of coastal aquaculture in the Indo-Pacific Region. In: *Coastal Aquaculture in the Indo-Pacific Region*. (Ed. T. V. R. Pillay) F. A. O. Fishing News (Books) Ltd., pp. 2-25.
- LINNAEUS, C. 1754. *Genera plantarum*. Holmiae. 500 pp.
- MAHONTY, C. B. 1956. Fishery byproducts industry in India—Seaweeds. in : *Progress of Fisheries Development in India-Cuttack*.
- MAIRH, O. P. 1982. Seasonal variation in alginic acid and viscosity of sodium alginate from a brown alga *Cystoseira indica* (Thivy et Doshi) Mairh, from Port Okha *Seaweed Res Util.*, 2 (1) : 43-46.
- MAIRH, O. P. AND V. KRISHNAMURTHY. 1988. Observations on the germination of spores and growth of germlings in a *Cystoseira*. *Jour. Indian bot. Soc.* 47 : 256-263.
- MAIRH, O. P. AND P. SREENIVASS RAO. 1978. Culture studies on *Gelidium pusillum* (Stack). *Le Jolis. Bot. Mar.*, 21 (3) : 169-174.
- MAIRH, O. P., P. C. THOMAS, B. K. RAMAVAT AND P. SREENIVASA RAO. 1979. Fertilizer pellets and their application in the field cultivation of *Gelidiella acerosa* (Forssk.) Feld et Hamel. *Proc. Int. Symp. Marine Algae of Indian Ocean Region*, CSMCRI, Bhavnagar, India. Abstract No. 12.
- MATHIESON, A. C. 1969. The promise of seaweed. *Oceanology Intl.*, Jan/Eeb. 1969. pp. 37-39.
- MEHTA, B. R. AND R. G. PAREKH. 1978. Mannitol content in brown algae of the coast of Sourashtra. *Bot. Mar*, 21 (4) : 251-252.



- MEHTA, V. C., B. S. TRIVEDI, K. K. BOKIL AND M. R. NARAYANA. 1967. Seaweeds as manure: I- Studies on nitrification. *Proc. Semi. Sea Salt and Plants*, CSMCRI, Bhavnagar. pp. 357-365.
- MICHANEK, G. 1975. Seaweed resources of the Ocean, F. A. O. *Fish. Tech. Rep.* No. 138. pp 1-126.
- MISRA, J. N. 1966. *Phaeophyceae in India*. I. C. A. R. New Delhi, pp. 1-203.
- MITRA, G. 1946. *Development of Chilka Lake, Cuttack*
- MIURA, A. 1975. *Porphyra* cultivation in Japan. In : *Advance of phycology in Japan*. (Ed. J. Tokida AND H. Hirose) Dr. W. Jung b. v. Publishers, The Hague. pp. 273-304.
- MOHAN JOSEPH, M. AND V. KRISHNAMURTHY. 1977. Studies on the shedding of carpospores in *Gracilaria corticata* J. Ag. *Seaweeds Res Util.*, 2 (1) : 1-8.
- MURTHY, M. S. AND P. RADIA. 1978. Eco-biochemical studies on some economically important intertidal algae from Port Okha (India). *Bot. Mar.*, 21 (7) : 417-422.
- NAQVI, S. W. A., MITTAL, S. Y. KAMAT, SOLIMABI AND C. V. G. REDDY. 1979. Bromine content in some seaweeds of Goa (central west coast of India). *Bot. Mar.*, 22 (7) : 455-457.
- NAQVI, S. W. A., SOLIMABI, S. Y. KAMAT, L. FERNANDES AND C. V. G. REDDY. 1981. Screening of some marine plants from the Indian coast for biological activity. *Bot. Mar.*, 24 (1) : 51-55.
- NEELA, M. V. 1956. Analysis of seaweeds. *Home. Sci. Bull. Women's Christian Coll.*, Madras.
- NEISH, I. C. 1979. Developments in the culture of algae and seaweeds and the future of the industry. In : *Advances in Aquaculture* (Ed. T. V. R. Pillay and Wm. A. Dill) FAO Fishing News (Books) Ltd. England pp. 395-402.
- OZA, R. M. 1971. Effect of IAA on the growth of fragment of *Gracilaria corticata*, J. Ag. *Seaweed Res. Util.*, 1 : 48-49.
- OZA, R. M. 1978. Studies on Indian *Gracilaria*. V. Seasonal variation in agar and gel strength of *Gracilaria corticata*. J. Ag. occurring on the coast of Veraval. *Bot. Mar.*, 21 (3) : 165-167.
- OZA, R. M. AND V. KRISHNAMURTHY. 1968. Studies on carposporic rhythm of *Gracilaria verrucosa* (Huds.) Papenf. *Bot. Mar.*, 11 (1-4) : 118-121.
- OZA, R. M. AND P. SREENIVASA RAO. 1977. Effect of different culture media on growth and sporulation of laboratory raised germlings of *Ulva fasciata* *Bot. Mar.*, 20 (7):427-431.
- PAPENFUSS, G. F. 1955. Classification of the algae. In: *A Century of Progress in the Natural Sciences, 1853-1953*. Calif. Acad. Sci. San Francisco. pp. 115-224.
- PAREKH, R. G. AND VISWESWARA RAO. 1964. Extraction of bulk proteins from the green seaweed, *Ulva rigida*. *Indian Jour. Tech.*, 2: 387.
- PAREKH, R. G., L. V. MARU AND M. J. DAVE. 1977. Chemical composition of green seaweeds of Saurashtra Coast. *Bot. Mar.* 20 (6): 359-362.
- PARIJA, P AND B. PARIJA. 1946. Algal succession on a rocky island named Charai Guha in the Chilka Lake. *Jour. Indian bot. Soc.* (M. O. P. Iyengar Commemo. Vol.) pp. 375-379.

- PARKER, H. S. 1974. The culture of red algal genus *Eucheuma* in the Philippines *Aquaculture*, 3 (4): 425-439.
- PASCHER, A. 1914. *Ber. dtsh. bot. Ges.* 32, 136.
- PATEL, B. A. AND G. V. JOSHI. 1967. Seasonal variations in chemical composition in *Ulva lactuca* and seawater. *Indian Jour. exp. Biol.*, 5: 236-238.
- PATEL, J. B., B. V. GOPAL, V. R. NAGULAN, K. SUBBARAMAIAH AND P. C. THOMAS. 1979. Experimental field cultivation of *Gelidiella acerosa* at Ervadi, India. *Proc. Int Symp. Marine Algae of the Indian Ocean Region*, CSMCRI, Bhavnagar, India. pp. 24-25 (Abstract).
- PATEL, J. B., B. V. GOPAL, V. R. NAGULAN, K. SUBBARAMAIAH AND P. C. THOMAS, 1980. Experimental field cultivation of *Gelidiella acerosa* at Ervadi, India. *Symp. Coastal Aquaculture*, Marine Biological Association of India, Cochin p. 189 (Abstract).
- PERCIVAL, E. 1968. Marine algal carbohydrates. *Oceanogr. Mar. biol. Ann. Rev.*, 6: 137-161.
- PILLAI, V. K. 1955 a. Observations on the ionic composition of bluegreen algae growing in saline lagoons. *Proc. natn. Inst. Sci. India*, 21: 90-102.
- PILLAI, V. K. 1955 b. Utilization of natural byproducts for the cultivation of blue-green algae. *Curr. Sci.*, 24: 21.
- PILLAI, V. K. 1955 c. Water soluble constituents of *Gracilaria lichenoides*. *Jour. Sci. Indust. Res. (India)*, 14 B. 473-477.
- PILLAI, V. K. 1956. Chemical studies on Indian seaweeds. I: Mineral constituents. *Proc. Indian Acad. Sci.*, B. 44: 3-29.
- PILLAI, V. K. 1957 a. Chemical studies on Indian seaweeds. II: Partition of Nitrogen. *Proc. Indian Acad. Sci.*, B 45: 43-63.
- PILLAI, V. K. 1957 b. Chemical studies on Indian seaweeds. III: Partition of Sulphur. *Proc. Indian Acad. Sci.*, B 45: 101-121.
- PILLAI, V. K. 1957 c. Alginic acid from *Sargassum* seaweeds. *Res. Ind.* 2: 70-71.
- PILLAI, V. K. 1964. Studies on the use of alginates in frozen fishery products. *Fishery. Tech.* 1: 176-179.
- PRESCOTT, G. W. 1968. *The algae: A review*. Houghton Mifflin Co., Boston 436 pp.
- PROVASOLI, L. 1957. Effect of plant hormones on *Ulva*, *Biol. Bull.*, 114: 375-384.
- QASIM, S. Z. AND M. V. M. WAFAR. 1979. Occurrence of living corals at several places along the west coast of India. *Mahasagar*. 12 (1): 53-58.
- RAGOTHAMAN, G. 1979. Littoral algal survey of south Gujarat coast (Devka, Golvad and Daman). *Proc. Int. Symp. Marine Algae of the Indian Ocean Region* CSMCRI, Bhavnagar, India. p. 6 (Abstract).

- RAJU, P. V. 1971. The effect of in situ application of growth hormones and fertilizers on photosynthetic  $c^{14}$  incorporation in some marine algae. *Bot. Mar.* 14 (2): 129-131.
- RAJU, P. V. AND P. C. THOMAS. 1971. Experimental field cultivation of *Gracilaria edulis* (Gmel.) Silva. *Bot. Mar.*, 14 (2): 71-75.
- RAJU, P. V. AND R. VENUGOPAL. 1971. Appearance and growth of *Sargassum plagiophyllum* (Mert.) C. Ag. on a fresh substratum. *Bot. Mar.* 14 (1): 36-38.
- RAMA RAO, K. 1970. Studies on growth cycle and phycocolloid content in *Hypnea musciformis* (Wulf.) Lamour. *Bot. Mar.*, 13 (2): 163-165.
- RAMA RAO, K. 1979. Studies on Indian Hypneaceae—V. Spores and natural propagules in the selected species of *Hypnea*, the potential Indian Carrageenophyte, for its field cultivation. *Proc. Int. Symp. Marine Algae of the Indian Ocean Region*. CSMCRI, Bhavnagar, India. pp 30-31 (Abstract).
- RAMA RAO, K AND V. KRISHNAMURTHY. 1968. Study of the preparation and properties of phycocolloid from *Hypnea musciformis* (Wulf.) Lamour. from Veraval, Gujarat coast, *Bot. Mar.*, 11: 129-133.
- RAMA RAO, K. AND P. C. THOMAS. 1974. Shedding of carpospores in *Gracilaria edulis* (Gmel.) Silva. *Phykos*, 13 (1) : 54-59.
- RAMA RAO, K. AND V. KRISHNAMURTHY. 1978. Studies on Indian Hypneaceae. I. Seasonal variation in phycocolloid content in two species of *Hypnea* (Gigartinales, Rhodophyceae). *Bot. Mar.*, 21 (4): 257-259.
- RAMA RAO, K. AND K. SUBBARAMAIAH. 1980. A technique for the field cultivation of *Hypnea musciformis* (Wulf.) Lamour., a carrageenophyte. *Symp. Coastal Aquaculture*. M. B. A. I., Cochin, India. p. 189 (Abstract).
- RANDHAWA, M. S. 1930. Historical review. Address; in: *Proc. Symp. Algology*, ICAR, New Delhi, pp. 4-24.
- ROUND, F. E. 1973. *The Biology of the Algae*, Edward Arnold, London-278 pp.
- RYTHER, J. H. 1968 a. *Porphyra* (Nori) culture in Japan. The status and potential of aquaculture particularly invertebrate and algae culture. Part II. Invertebrate and algae culture. (Reproduced by National Technical Service, Springfield, Va. 22151). pp. 228-241.
- RYTHER, J. H. 1968 b. *Undaria* culture in Japan. The status and potential of aquaculture; particularly invertebrate and algae culture. Part II. Invertebrate and algae culture (Reproduced by National Technical Information Service, Springfield, Va. 22151). pp. 242-248.
- SADASIVAN PILLAI, K. 1961. Alginic acid from *Sargassum* seaweeds of Indian coasts—Its extraction on a cottage industry basis. *Chemical age of India*, 12 : 425-430.
- SADASIVAN PILLAI, K. AND N. S. VARIER. 1952. Studies on the structure of alginic acid from the *Sargassum* seaweeds of Cape Comorin. *Jour. Proc. Inst. Chem.*, (India), 24 : 205.
- SAITO, Y. 1975. *Undaria*. In : *Advance of Phycology in Japan* (Ed. J. Tokida and H. Hirose) Dr. W. Jung b. v. Publishers, The Hague. pp. 304-320.

- SAITO, Y. 1979. Seaweed aquaculture in the North West Pacific. In : *Advance in Aquaculture* (Ed. T. V. R. Pillay and Wm. A. Dill). F. A. O. Fishing News (Books) Ltd. England. pp. 402-410.
- SARMA, Y. S. R. K. AND M. KHAN. 1980. Algal taxonomy in India. Botanical records and monographs-2. Today and Tomorrow's Printers and Publishers, New Delhi. pp. 1-153.
- SHAH, H. N. AND A. V. RAO. 1969. Recovery of mannitol from Indian brown seaweeds. *Res. Ind.*, 14 (3) : 117-119.
- SHAH, H. N., I. A. Mody and A. VISWESWARA RAO. 1967. Seasonal variation of viscosity of sodium alginate from *Sargassum* species and the preparation of high viscosity alginates, *Indian Jour. Tech.*, 5 : 269-270.
- SITAKARA RAO, V. AND U. K. TIPNIS. 1964. Protein content of marine algae from Gujarat coast. *Curr. Sci.*, 33 : 16-17.
- SITAKARA RAO, V. AND U. K. TIPNIS 1967. Chemical composition of marine algae from Gujarat coast. *Proc. Semi. Sea Salt and Plants*, CSMCRI, Bhavnagar, pp. 277-288.
- SMITH, G. A. 1955. *Cryptogamic Botany*; Vol. I. Mc Graw Hill Book Co., New York. 546 pp.
- SOLIMABI AND S. W. A. NAQVI. 1976. Alginic acid content of some brown seaweeds of Goa. *Mahasagar* 8 (1 & 2) : 97-99.
- SOLIMABI AND B. DAS. 1977. Distribution of iodine in marine algae of Goa region. *Indian J. mar. Sci.*, 6 (2) : 180-181.
- SOLIMABI, B. DAS, S. Y. KAMAT, L. FERNANDES AND C. V. G. REDDY. 1980. Seasonal changes in carrageenan and other biochemical constituents of *Hypnea musciformis*. *Indian. J. mar. Sci.*, 9 (2): 134-136.
- SOLIMABI, B. DAS, P. K. MITTAL AND S. Y. KAMAT, 1981. Bromine and iodine contents in sponges and algae of the Andaman Sea. *Indian J. mar. Sci.*, 10 (3): 301-302.
- SREENIVASA RAO, P. 1967. Laver cultivation in Japan. *Salt Res. Ind.*, 4 (4): 141-144.
- SREENIVASA RAO, P. 1969. Systematics, ecology and life history of Indian Gelidiales with special reference to agarophyte *Gelidiella acerosa* (Forsskal) Feldman et Hamel. *Salt Res. Ind.*, 6: 46-47.
- SREENIVASA RAO, P. 1970. Systematics of Indian Gelidiales. *Phykos*. 9: 63-78.
- SREENIVASA RAO, P. AND S. R. KALE. 1969 Marine algae from a little known place of Gujarat coast I. Algae from Gopnath. *Phykos*, 8: 71-82.
- SREENIVASA RAO, P. AND Y. A. Shelat. 1979. Antifungal activity of Indian seaweed extracts. *Proc. Int. Symp. Marine Algae of the Indian Ocean Region*. CSMCRI, Bhavnagar, India. p. 47 (Abstract).
- SREENIVASA RAO, P. AND K. S. PAREKH. 1981. Antibacterial activity of Indian seaweed extracts. *Bot Mar.*, 14 (11): 577-582.
- SREENIVASA RAO, P., E. R. R. IYENGAR AND F. THIVY. 1964. Survey of algin bearing seaweeds at Adatra reef, Okha. *Curr. Sci* 33: 464-465.

- SREENIVASA RAO, P., H. H. PAREKH, B. K. RAMAVAT AND S. B. BHATT. 1979 a. Preparation and properties of liquid seaweed fertilizer. *Proc. Int. Symp. Marine Algae of the Indian Ocean region*. CSMCRI, Bhavnagar, India p. 57 (Abstract).
- SREENIVASA RAO, P., S. J. TARWADE, K. S. R. SARMA, K. ANJANEYULU AND H. M. MODY, 1979 b. Seaweed as a source of energy: Production of fuel gas from seaweed, *Sargassum*. *Proc. Int. Symp. Marine Algae of the Indian Ocean Region*, CSMCRI, Bhavnagar, India. p. 56 (Abstract).
- SREENIVASA RAO, P., K. S. PAREKH AND H. H. PAREKH. 1979 c. Antibacterial activity of different fractions of seaweed extracts. *Proc. Int. Symp. Marine Algae of the Indian Ocean Region*, CSMCRI, Bhavnagar, India. p. 47 (Abstract).
- SREENIVASA RAO, P., K. S. PAREKH, H. H. PAREKH, S. B. TRIVEDI AND B. A. DAVE. 1979 d. Effect of seaweed extracts on *Mycobacterium tuberculosis*. *Proc. Int. Symp. Marine Algae of the Indian Ocean Region*, CSMCRI, Bhavnagar, India. p. 47 (Abstract).
- SRINIVASAN, K. S. 1946. Ecology and seasonal succession of the marine algae at Mahabalipuram (Seven Pagodas) near Madras. *Jour. Indian. bot. Soc.*, (M. O. P. Iyengar commemo. Vol.) pp. 267-278.
- SRINIVASAN, K. S. 1950. Distribution patterns of marine algae in Indian seas *Proc. Symp. Algology*, ICAR, New Delhi. pp. 219-242.
- SRINIVASAN, K. S. 1965. Indian botany in retrospect with particular reference to algal systematics. *Jour. Asiatic Soc., Bengal*, 7: 49-78.
- SRINIVASAN, K. S. 1966. Conspectus of *Sargassum* species from Indian territorial waters. *phykos*, 5: 127-129.
- SRINIVASAN, R. AND T. SANTHANARAJA. 1967. Studies on the extraction and properties of agar-agar from the seaweed *Gracilaria* species in Madras State. *Madras Jour. Fish.*, 3: 146-151.
- STANFORD, E. C. C. 1883. *Chem. News.*, 47: 254-257, 262-269.
- STANIER, R. Y. AND VAN NIEL, C. B. 1962. *Bact. Rev.*, 42, 17.
- SUBBARAMAIAH, K. 1937. Ascorbic acid content and growth in *Ulva fasciata* Delile, *Phykos*, 6: 115-117.
- SUBBARAMAIAH, K. 1970. Growth and reproduction of *Ulva fasciata* Delile in nature and in culture. *Bot. Mar.*, 13 (1): 25-27.
- SUBBARAMAIAH, K., AND V. KRISHNAMURTHY. 1967. Laboratory culture of seaweeds. *Proc. Semi. Sea Salt and Plants*, CSMCRI, Bhavnagar pp. 321-326.
- SUBBARAMAIAH, K., S. R. KALE AND V. KRISHNAMURTHY. 1967. Gametes and germings of *Ulva fasciata* Delile. *Curr. Sci.*, 36: 128-129.
- SUBBARAMAIAH, K., K. RAMA RAO, P. C. THOMAS, M. R. P. NAIR, B. V. GOPAL AND V. R. NAGULAN. 1975. Cultivation of *Gelidiella acerosa*. *Salt Res. Ind.*, 11 (1): 33-36.

- SUBBARAMAIAH, K., K. RAMA RAO, M. R. P. NAIR, C. V. S. KRISHNAMURTHY AND M. PARAMASIVAM 1979 a. Marine algal resources of Tamil Nadu. *Proc. Int. Symp. Marine Algae of the Indian Ocean Region*, CSMCRI, Bhavanagar, India. p. 14 (Abstract.)
- SUBBARAMAIAH, K., K. RAMA RAO AND M. R. P. NAIR. 1979 b. Marine algal resources of Lakshadweep. *Proc. Int. Symp. Marine Algae of the Indian Ocean Region.*, CSMCRI, Bhavnagar, India. pp. 6-7 (Abstract).
- SUBBA RAO, G. N. 1965. Use of seaweeds directly as human food. *Indo-Pacific Fish. Coun. Reg. Studies. No. 2*: 1-32.
- SUBBA RAO, P. V., K. RAMA RAO AND K. SUBBARAMAIAH. 1977. Screening of certain red seaweeds for phycocolloids. *Seaweed Res. Util.*, 2 (2): 82-86
- SUBRAHMANYAN, R. 1967. Methods of assessing seaweed resources and problems. *Proc. Semi. Sea Salt and Plants*, CSMCRI, Bhavnagar. pp. 311-314.
- SUMITRA VIJAYARAGHAVAN, M. D. RAJAGOPAL AND M. V. M. WAFAR. 1980. Seasonal variations in biochemical composition of seaweeds from Goa coast. *Indian J. mar. Sci.* 9 (1): 61-63.
- TAMIYA, H. 1960. Role of algae as food. *Proc. Symp. Algology*, ICAR, New Delhi, pp. 379-389
- TAYLOR, W. R. 1964. The genus *Turbinaria* in eastern seas. *Jour. Linn. Soc. London (Botany)*, 58: 475-490.
- TEWARI A. 1975. The effect of a morphactin on the vegetative growth of *Gelidiella acerosa* *Phykos* 14 (1 & 2) : 125-128.
- TEWARI, A., M. PRASADA RAO AND V. KRISHNAMURTHY. 1968. Chemical composition of a species of *Porphyra* from Visakhapatnam. S. India. *Curr. Sci.*, 37: 138.
- THIVY, F. 1952. Investigations of seaweed products in India with a note on properties of various Indian agars. *Proc. Indo-Paci. Fish. Council, Sec. 2* : 173-175.
- THIVY, F. 1958. Economic seaweeds. In *Fisheries of West Coast of India*, Bangalore. pp. 74-80.
- THIVY, F. 1960. Seaweed utilization in India. *Proc. Symp. Algology*, ICAR, New Delhi, pp. 345-365.
- THIVY F. 1964. Marine algal cultivation. *Salt Res Ind.*, 1 (1) . 23-28.
- THOMAS, P. C 1977. Seasonal variation in the yield and physical properties of agar-agar from *Gracilaria verrucosa* (Hudson) Papenfuss. *Seaweed Res. Util.*, 2 (2) : 78-81.
- THOMAS, P. C. AND V. KRISHNAMURTHY. 1976. Agar from cultured *Gracilaria edulis* (Gmel.) Silva. *Bot. Mar.*, 19 : 115-117.
- THOMAS, P. C., K. RAMA RAO AND K. SUBBARAMAIAH 1975. Periodicity in growth and production of agar of *Gelidiella acerosa* (Forssk.) Feldman et Hamel. *Indian J. Mar. Sci.*, 4 (2) : 210-212.

- UMAMAHESWARA RAO, M. 1969 a. Catalogue of marine algae in the reference collection of the Central Marine Fisheries Research Institute. *Bull. cent. mar. Fish. Res. Inst.*, 9 : 37-48.
- UMAMAHESWARA RAO, M. 1969 b. Agar and algin yielding seaweeds of India. *Proc. 6th Int. Seaweed Symp.*, pp. 715-721.
- UMAMAHESWARA RAO, M. 1969 c. Seasonal variations in growth, alginic acid and mannitol contents of *Sargassum wightii* and *Turbinaria conoides* from the Gulf of Mannar, India. *Proc. 6th Int. Seaweed Symp.* pp. 579-584.
- UMAMAHESWARA RAO, M. 1970. The economic seaweeds of India. *Bull. cent. mar. Fish. Res. Inst. No. 20* : pp. 1-68.
- UMAMAHESWARA RAO, M. 1972 a. Coralreef flora of Gulf of Mannar and Palk Bay. *Proc. Symp. Corals and Coral Reefs* (1969). pp. 217-230.
- UMAMAHESWARA RAO, M. 1972 b. On the Gracilariaceae of the seas around India. *Jour. mar. biol. Ass. India*, 14 (2) : 671-696.
- UMAMAHESWARA RAO, M. 1972 c. Ecological observations on some intertidal algae of Mandapam coast. *Proc. Indian natl. Sci. Acad.*, 38 B (3 & 4) : 298-307.
- UMAMAHESWARA RAO, M. 1973. The seaweed potential of the seas around India. *Proc. Symp. on Living Resources of the Seas Around India* (1968). pp. 687-692.
- UMAMAHESWARA RAO, M. 1974. a. On the cultivation of *Gracilaria edulis* in the nearshore areas around Mandapam. *Curr. Sci.*, 43 (20) : 660-661.
- UMAMAHESWARA RAO, M. 1974 b. Observations on fruiting cycle, spore output and germination of tetraspores of *Gelidiella acerosa* in the Gulf of Mannar. *Bot Mar.*, 17 (4) : 204-207.
- UMAMAHESWARA RAO, M. 1976. Spore liberation in *Gracilaria corticata* J. Agardh growing at Mandapam. *J. exp. Mar. Biol. Ecol.*, 21 : 91-98.
- UMAMAHESWARA RAO, M. 1978. Seaweed resources of Andhra Pradesh. *Seaweed Res. Util.*, 3 (1 & 2) : 51-55.
- UMAMAHESWARA RAO, M. AND T. SREERAMULU. 1963. Vertical zonation and seasonal and variation in the growth of *Porphyra* on Visakhapatnam coast. *Curr. Sci.*, 32 : 173-174.
- UMAMAHESWARA RAO, M. AND T. SREERAMULU. 1970. An annotated list of the marine algae of Visakhapatnam (India). *Bot. Jour. Linn. Soc.*, 63 : 23-45.
- UMAMAHESWARA RAO, M. AND S. KALIMUTHU. 1972. Changes in mannitol and alginic acid contents of *Turbinaria ornata* (Turner) J. Agardh in relation to growth and fruiting. *Bot Mar.*, 15 : 57-59.
- UMAMAHESWARA RAO, M. AND N. KALIAPERUMAL. 1976. Some observations on the liberation and viability of oospores in *Sargassum wightii* (Greville) J. Ag. *Indian J. Fish.*, 23 (1 & 2), 232-235.

- UNNI, C. K. 1967. Natural radioactivity of marine algae. *Proc. Semi. Sea Salt and Plants*, CSMCRI: Bhavnagar. pp. 265-273.
- UNTAWALE, A. G. AND V. K. DHARGALKAR. 1975. *Report on the seaweed resources of the Goa coast*. N. I. O., Dona Paula, Goa. pp. 1-10.
- UNTAWALE, A. G., N. B. BHOSLE AND V. K. DHARGALKAR. 1977. Properties of phycocolloid extraction from seaweeds of Goa. *Indian J. mar. Sci.*, 6 (2) : 181-183.
- UNTAWALE, A. G., V. K. DHARGALKAR, V. V. AGADI AND T. G. JAGTAP. 1979. Marine algal resources of the Maharashtra coast. *Tech. Report*. Natl. Inst. of Oceanography, Goa. 48 pp.
- UNTAWALE, A. G., V. K. DHARGALKAR AND V. V. AGADI. 1983. *List of marine algae from India*. N. I. O., Dona Paula, Goa. pp. 1-42.
- VALSON, A. P. 1955. Alginic acid content of some of the common seaweeds of the Gulf of Mannar area. *Curr. Sci.*, 24: 343-345.
- VARIER, N. S. AND K. SADASIVAN PILLAI. 1952. Mannitol from *Sargassum* seaweeds. II. Optimum conditions for extraction of alginic acid from *Sargassum* seaweeds of Cape Comorin. *Bull. cent. Res. Inst.*, 2: 39.
- VARMA, R. P. 1960. Flora of the pearl beds off Tuticorin. *Jour. mar. biol. Ass. India*, 2: 221-225.
- VARMA, R. P. AND K. KRISHNA RAO. 1962. Algal resources of Pamban area. *Indian J. Fish.* 9: 205-211.
- VILHELM, J. 1931. *Archaeophyta and Algophyta*. Prague.
- VINOGRADOV, A. P. 1953. The elementary chemical composition of marine organisms. *Sears, Foundation for Marine Res.*, No 11.
- VISWESWARA RAO, A. 1964. Protein from *Ulva*. *Salt Res. Ind.*, 1: 37.
- VISWESWARA RAO, A. AND I. C. MODY. 1964. Extraction of alginic acid and alginates from brown seaweeds. *Indian Jour. Tech.*, 3 (8): 261-262.
- VISWESWARA RAO, A. K. N. PATEL AND H. N. SHAH. 1965. Manufacture of agar-agar from red seaweeds. *Res. Ind.*, 10: 131-133.
- WHITTAKER, R. H. 1969. *Science*, N. Y., 163, 150.
- YAPHE, W. 1959. The determination of kappa carrageenan as a factor in the classification of Rhodophyceae. *Canad. Jour. Bot.*, 37: 751-757.
- ZINGDE, M. D., S. Y. S. SINGBAL, C. P. MORAES AND C. V. G. REDDY. 1976. Arsenic, Copper Zinc and Manganese in the marine flora and fauna of coastal and estuarine waters around Goa. *Indian J. mar. Sci.*, 5: 212-217.